

# Administrarea subcutanata a medicamentelor in ingrijirea paliativa



Luciana Negulescu  
Asistent medical  
Asociatia pentru Servicii Mobile de Ingrijire Paliativa  
Bucuresti

# Administrarea subcutanata a medicamentelor

## Indicatii:

- tumori sfera ORL,
- varsaturi,
- disfagie,
- stare terminala,
- coma.



# Administrarea subcutanata a medicamentelor

## Avantaje:

- evitarea administrării IV, IM (durere, spargerea venei)
- frecvența redusă a intepaturilor,
- bun control simptomatic,
- fixarea acului (functional 5-7 zile),
- confort crescut pentru pacient inclusiv la domiciliu.



# Administrarea subcutanata a medicamentelor

## Dezavantaje:

- nu toate medicamentele pot fi administrate SC (substante uleioase sau iritante),
- maxim 2-2,5 ml la o administrare.

## Incidente (rare):

- posibila durere la locul injectiei,
- sangerare,
- infundarea acului,
- iritatie locala\*.



# Administrarea subcutanata a medicamentelor

## Medicamentele care pot fi administrate SC:

Antialgice: ketorolac  
tramadol  
morfina



# Administrarea subcutanata a medicamentelor

## Medicamentele care pot fi administrate SC:

Antiemetice: metoclopramid  
ondansetron  
scobutil



# Administrarea subcutanata a medicamentelor

## Medicamentele care pot fi administrate SC:

Altele:

dexametazona

haloperidol

midazolam

levomepromazina

fenobarbital


ranitidina

atropina



# Administrarea subcutanata a medicamentelor

## Zone ale administrarii SC

















































# Administrarea subcutanata a medicamentelor

## Materiale folosite la administrarea SC:

- microperfuzoare (fluturasi) cu dimensiunea de 25 G,
- siringa 2 ml,
- pansament ocluziv transparent “tegaderm”, pentru fixare ac
- tampon cu alcool.



# Compatibilitatea substantelor in siringa

	Morfina	Midazo -lam	Haloperi -dol	Scobu -til	Dexame -tazona	Metoclo -pramid	Trama -dol	Raniti -dina
Morfina							—	
Midazo -lam								
Haloperi -dol								?
Scobutil								
Dexame -tazona								?
Metoclo -pramid								?
Trama -dol	—							?
Raniti -dina			?		?	?	?	

# Bibliografie

Ferrel B, Coyle N, Textbook of Palliative Nursing, 2<sup>nd</sup> Edition Oxford University Press, 2007

Negro S, Azuara ML, Sanchez Y, Reyes R, Barcia E, *Physical compatibility and in vivo evaluation of drug mixture for subcutaneous infusion to cancer patients in palliative care*, Supportive Care in Cancer, 2002

Neuenschwander H et al, *Medicine palliative*, Ligue suisse contre le cancer, 2007

